ESA 1996 Abstract Page 1

Author to contact: Kyle McDonald

Institution: Jet Propulsion Laboratory

Mailing address:
Mail Stop 300-233
Jet. Propulsion Laboratory
4800 Oak Grove Drive
Pasadena, CA 91109

Phone number: 818-354-3263

Finail address: kyle.mcdonald@jpl.nasa.gcv

Presenter: Kyle McDonald

Preference: oral

Presenter is a non-member

1st Choice: 63 2nd Choice: 38

(Special session on the BORFAS experiment, if one is being organized)

MCDONALD, KYLE, JOBEA WAY, REINER ZIEMER AND and ERIC RIGNOT. Jet Propulsion laboratory, 4800 (a k Grove orive, Pasadena, CA 91109, USA. Application of imaging radar tomonite graving season duration at BOREAS.

Improved estimates of the seasonal duration of carbon exchange processes in boreal regions may improve corrent entiretes of net annual CO@2@ flux at high latitudes. For coniferous tree species, the summer frost-free period bounds growing season length and period of carbon intake. For the broader landscape, the spring soil than bounds the period of soil respiration and carbon release. In address ag the use of imaging radar as a tool for estimating periods of carbon exchange, one must understand (1) the relationship between canopy freeze thaw state and the period of photosynthetic and respiration activity and (2) the sensitivity of radar to vegetation and soil freeze/thaw processes. We deployed several suites of measurement systems to monitor in situ soil and vegetation temperature, xylem sap flux, vegetation dielectric constant, and within-canopy meteorologic variables at the Boreal Ecosystem Atmosphere Study (BOREAS) in Canada. Imaging radar observations were obtained with the ERS-1 SAR throughout 1994. These data were supplemented with CO@2@ flux determined from models and tower-base | measurements. Results show that changes in radar backscatter were vel-correlated with the start of both soil respiration and vegetation COCCO intake, and demonstrate radar's utility for providing useful input to CO@2@ flux estimates.